arXiv:1609.08683v1 [astro-ph.SR] 27 Sep 2016

On the formation of DA white dwarfs with low hydrogen contents: Preliminary Results.

Miller Bertolami, M. M.[1], Althaus, L. G.[1,2], and Córsico, A. H.[1,2]

[1]*Instituto de Astrofísica La Plata, CONICET-UNLP, Paseo del Bosque s/n, (B1900FWA) La Plata, Argentina;* mmiller@fcaglp.unlp.edu.ar, althaus@fcaglp.unlp.edu.ar, acorsico@fcaglp.unlp.edu.ar

[2]*Facultad de Ciencias Astronómicas y Geofísicas, Universidad Nacional de La Plata, Paseo del Bosque s/n, (B1900FWA) La Plata, Argentina.*

Abstract. Systematic photometric and asteroseismological studies in the last decade support the belief that white dwarfs in the solar neighborhood harbor a broad range of hydrogen-layer contents. The reasons behind this spread of hydrogen-layer masses are not understood and usually misunderstood. In this work we present, and review, the different mechanisms that can (or cannot) lead to the formation of white dwarfs with a broad range hydrogen contents.

1. Introduction

White dwarf stars (WD) are the most common end point of stellar evolution (Althaus et al. 2010). In the most simple scenario, WDs are formed when the progenitor stars lose their external envelopes at the end of the Asymptotic Giant Branch (AGB) and contract at constant luminosity to the WD cooling sequence. Within this simple picture stellar evolution theory usually predicts the formation of WD with pure hydrogen (H) atmospheres and with a total H content of about $M_{\rm H}^{\rm WD} \sim 10^{-3}$–$10^{-5} M_{\odot}$. About 80% of the spectroscopically identified WDs are characterized by H-rich atmospheres (usually referred as DA WDs). Considerable progress in the study of WDs has been possible because some of these stars pulsate. The study of the pulsational pattern of variable DA WDs (DAVs) through asteroseismology has provided valuable constraints to their mechanical properties, such as the core composition, the outer layer chemical stratification and the stellar mass. Early asteroseismological fittings suggested a range of H layer masses between 10^{-4} and 10^{-10} $M_{\odot}$ —see Bradley (2001) and references therein. The mass of the outer H layer is an important point regarding the spectral evolution theory —see Bergeron et al. (2011), and also Rolland et al. in these proceedings). Tremblay & Bergeron (2008) presented a detailed photometric study of the spectral evolution of cool WDs ($T_{\rm eff} < 15000$K). They determined the ratio of helium(He)- to H-atmosphere WDs as a function of temperature and concluded that $\sim$ 15% of the DA WD between 15000 and 10000 K will be transformed into He-atmosphere, non-DA, WD at lower temperatures. From a model of convective mixing these authors then concluded that 85% of DA WD harbor H-layer masses larger than $M_{\rm H}^{\rm WD} \sim 10^{-6} M_{\odot}$, with the remaining 15% harboring less massive H layers. The existence a range of H-layer masses was later confirmed by the systematic studies of Castanheira & Kepler (2009) and Romero

et al. (2012) by asteroseismological means. In particular, the study of Romero et al. (2012), finds 77% of their DAV sample (34 out of 44) to harbor H-layer masses larger than $M_{\rm H}^{\rm WD} = 10^{-6} M_{\odot}$, in good agreement with the photometric determinations of Tremblay & Bergeron (2008).

In this context, it is interesting to analyze the evolutionary channels that can lead to low H contents in DA WDs. Already D'Antona & Mazzitelli (1990) reviewed several possible channels for the formation of DA WDs with low H contents. These channels were: a) The action of winds during the stationary He burning phase after a late thermal pulse (Schoenberner 1979; Iben 1984); b) a late thermal pulse during the WD cooling track, where the H envelope is violently burned (Iben 1984; Iben & MacDonald 1995); and c) a diffusion/self-induced nova in the WD cooling track (Iben & MacDonald 1986). The study of Miller Bertolami et al. (2011) showed that the diffusion-induced nova does not lead to a significant decrement of the H content of WDs. On the contrary scenarios a) and b), now called "late" and "very late" thermal pulses (LTP, and VLTP respectively; Blöcker 2001) have been proven to significantly reduce the H content of WDs. The quantitative exploration by Althaus et al. (2005) and Miller Bertolami et al. (2005) showed that H contents as low as $M_{\rm H}^{\rm WD} \sim 10^{-7} M_{\odot}$ could be produced in LTPs. VLTPs, on the other hand, have been usually understood to produce non-DA WDs (Iben 1984; Althaus et al. 2005; Werner & Herwig 2006). Although numerical simulations never predict the burning of the complete H content of the star (Miller Bertolami et al. 2006), it has been argued that whatever the traces of H that may be left by the VLTP they will very likely be shed off by mass loss during the subsequent giant phase — Werner & Herwig (2006). While this is a possibility it has to be emphasized that, once the star is back on its giant phase, H is diluted in the more massive convective envelope of the born again AGB star ($\sim 10^{-3} M_{\odot}$). The star would need to lose all that mass in order to get rid of its whole H content. In addition, the study of Miller Bertolami & Althaus (2007) suggested that the amount of H burned in low-mass VLTPs might be only a small fraction of the total H content of the star. Consequently, DA WDs with low H contents could be formed both through LTP and VLTP events.

In this work we present preliminary explorations aimed at a quantitative assessment of the range of WD with low H contents predicted by different post-AGB channels in the light of state-of-the-art stellar evolution models (Miller Bertolami 2016). Although it was already clear from early works that the H content of the future WD cannot be arbitrarily reduced by winds, during the AGB or H-burning phases (Paczyński 1971; Schoenberner 1979, 1987), some misconceptions have arisen in recent works. Consequently, before presenting the predictions of the LTP and VLTP channels we will discuss why enhanced mass loss on the AGB and/or in H-burning post-AGB sequences cannot lead to DA WDs with low H contents.

2. The misconceptions: Winds on the AGB and H-burning post-AGB phases.

Fig. 1 shows the dependence of the H content of the resulting WD ($M_H^{\rm WD}$) with the initial progenitor metallicity ($Z_{\rm ZAMS}$) and final mass ($M_{\rm WD}$) as predicted for H-burning post-AGB sequences by Miller Bertolami (2016). As it is apparent in Fig. 1 the $M_H^{\rm WD} - M_{\rm WD}$ relation of WD resulting from H-burning post-AGB sequences is not very sensitive to the value of $Z_{\rm ZAMS}$ and restricted to the range $10^{-3} M_{\odot} < M_{\rm H}^{\rm WD} < 10^{-5} M_{\odot}$. Consequently, WDs evolved from H-burning post-AGB objects show a rather tight $M_{\rm H}^{\rm WD} - M_{\rm WD}$ relation.



Figure 1. Solid Lines: $M_H^{\rm WD} - M_{\rm WD}$ relations of DA WDs resulting from H-burning post-AGB sequences with different $Z_{\rm ZAMS}$ (Miller Bertolami 2016). Black dots: $M_{\rm H}^{\rm WD} - M_{\rm WD}$ values of DA WDs constructed from an initially $M_{\rm ZAMS} = 1.25 M_\odot$, $Z = 0.001$ star by artificially increasing the AGB winds by a factor of 3 and 10.

As noted by Paczyński (1971), and also Schoenberner (1987), H-burning post-AGB models have a very tight relationship between the effective temperature ($T_{\rm eff}$) and the envelope mass of the post-AGB object. As long as the envelope can be considered in thermal equilibrium, the location of a H-burning post-AGB remnant on the HR-diagram is independent of the mass loss history and only dependent on the value of the envelope mass. As a consequence an enhancement in the post-AGB winds does speed up the post-AGB evolution but does not reduce the value of $M_H^{\rm WD}$. To check this argument we recomputed the sequence with $M_{\rm ZAMS} = 1.25 M_\odot$, $Z = 0.001$ presented by Miller Bertolami (2016) by artificially enhancing the winds on the post-AGB phase. Our computations show that even an enhancement of a factor 100 in the post-AGB winds affects neither the $T_{\rm eff}$-$M_{\rm H}$ relation nor the final H content of the WD ($M_H^{\rm WD}$).

The intensity of the winds of AGB stars are not directly relevant for the final H content of the WD. Fig. 1 shows the impact of increasing the AGB winds up to an order of magnitude. An enhancement of AGB winds leads to a shortening of the AGB. With less time for the H-free core to grow during the thermal pulses, enhancing the winds on the AGB leads to a smaller final mass ($M_{\rm WD}$) for the same initial mass ($M_{\rm ZAMS}$). Due to the tight $M_{\rm WD}$-$M_{\rm H}^{\rm WD}$ relation this implies that more intense winds on the AGB lead to the formation of WD with higher H contents. This is true also when looked at at the same value of the remnant mass ($M_{\rm WD}$). Models of the same final mass but shorter AGB lifetimes are less compact and luminous and have, consequently, larger post-AGB envelope masses (Bloecker 1995; Miller Bertolami 2016).

Different AGB winds can indirectly affect the H content of the WD by affecting the frequency of late thermal pulses. Depending on how AGB winds are modulated with the surface temperature, brightness and composition of the star, the occurrence of late thermal pulses can be favored or not. And, as shown in the next section, late thermal pulses do indeed reduce the H content of the future WD.

3. Late Flashers: A formation channel for DA WD with low H contents



Figure 2. Evolution of the H content ($M_{\rm H}$) of stars during LTP and VLTP events (dashed line) as compared with their canonical evolution in H-burning post-AGB sequences from the main sequence to the final WD value (dotted line).

As discussed in the previous sections the occurrence of late thermal pulses, or the action of winds during the He-burning phase after a late thermal pulse can lead to an effective depletion of the H content of the resulting WD stars. In order to quantitatively assess the final H content expected from the VLTP and LTP events we have recomputed some of the sequences presented in Miller Bertolami (2016), by and ad-hoc tuning of the mass loss during the departure from the AGB in order to obtain a late thermal pulse. In particular we have computed three LTP sequences corresponding to $(M_{\rm ZAMS}, M_{\rm WD}, Z_{\rm ZAMS}) = (1.25M_{\odot}, 0.5900M_{\odot}, 0.001)$, $(1.50M_{\odot}, 0.5961M_{\odot}, 0.001)$, $(2.50M_{\odot}, 0.6200M_{\odot}, 0.01)$ and two VLTP sequences corresponding to $(M_{\rm ZAMS}, M_{\rm WD}, Z_{\rm ZAMS}) = (1.25M_{\odot}, 0.5894M_{\odot}, 0.001)$, $(2.50M_{\odot}, 0.6194M_{\odot}, 0.01)$. In Fig 2 we show the typical evolution of the H content of post-AGB objects that undergo LTP and VLTP events.

As discussed in Althaus et al. (2005), and shown in Fig. 2, after a LTP event the star returns to its cool giant structure for a second time. If third dredge up develops[1] then H is diluted in the rather deep convective envelope of the reborn red giant and the surface becomes H deficient. Although no H is burnt in this phase, some H can be lost due to stellar winds. Once the star departs from the AGB for a second time, the envelope contracts and heats up. Then the H located in the inner parts of the star is burnt and the total H content the present simulations is reduced to $M_{\rm H} \sim 10^{-6}$–$10^{-7} M_{\odot}$ (see Fig. 3). It is worth noting that final H content of the WD could be higher if the LTPs would have taken place closer to the AGB —a scenario termed AGB final thermal pulse, see Blöcker (2001).

In the case of VLTP events, the H content of the models is also reduced by the violent H flash that follows the thermal pulse —see Miller Bertolami et al. (2006) for a very detailed description of the event. After the post-VLTP object returns to the AGB, the evolution proceeds as in the LTP case. The remaining H is diluted in the convective envelope of the reborn AGB star and, when the star departs from the AGB for a second time H is burned again in the inner parts of the envelope as it heats up. Interestingly, and in agreement with our previous works (Miller Bertolami & Althaus 2007) we find that the H flash only burns part of the available H in low-mass VLTPs ($M_{\rm WD} \lesssim 0.6 M_{\odot}$). This is because in low-mass VLTP events burning only a fraction of the available H injects enough energy in the envelope of the star to expand it to its giant dimensions. On the other hand, in more massive remnants undergoing VLTP events, burning the whole H content is not enough to expand the envelope and H burning only stops when the He-flash driven expansion finally pushes the star back to the AGB. As a consequence, in this preliminary explorations we find in VLTP events that the final H content of the star strongly depend on this mass threshold, being of the order of $M_{\rm H} \sim 10^{-11} M_{\odot}$ for the more massive sequence and much closer to that expected in LTP events for the less massive ones ($M_{\rm H} \sim 10^{-7.3} M_{\odot}$, see Figs. 2 and 3).

4. Discussion and future work

In this preliminary study we have shown that both flavors of late thermal pulses (VLTPs and LTPs) can lead to the formation of DA WDs with low H contents (see Fig. 3). In addition we have confirmed that uncertainties in AGB and post-AGB winds during the H-burning phase are not directly relevant for the formation of DA WDs with low H contents. In the case of LTPs our simulations predict values as low as $M_{\rm H}^{\rm WD} \sim 10^{-7} M_{\odot}$. As the amount of H remaining in this scenario is tightly connected to the the mass of the H-rich envelope at the moment of the He-flash, we expect a smooth transition from the canonical H-burning post-AGB value (Fig. 1) down to $M_{\rm H}^{\rm WD} \sim 10^{-7} M_{\odot}$. In addition, we have shown that current VLTP computations predict a bi-modal distribution for the value of $M_{\rm H}^{\rm WD}$. With VLTP remnants of masses $M_{\rm WD} \gtrsim 0.6 M_{\odot}$ resulting in $M_{\rm H}^{\rm WD} \sim 10^{-11} M_{\odot}$ and less massive VLTP remnants harboring H contents just below the LTP predictions ($M_{\rm H} \lesssim 10^{-7} M_{\odot}$). Future work will expand these preliminary computations and explore the role of both remnant masses and winds during the He-flash driven evolution.

[1]The occurrence of third dredge up depends on the initial mass of the model and on the number of thermal pulses (Miller Bertolami 2016).



Figure 3. H content of the WDs predicted by VLTP and LTP channels as compared with the canonical predictions of H-burning post-AGB sequences for two different initial metallicities.

Acknowledgments. Part of this work was supported by ANPCyT through grant PICT-2014-2708 from FonCyT and grant PIP 112-200801- 00940 from CONICET. M3B thanks the Alexander von Humboldt Foundation for a Return Fellowship. M3B and AHC thank the organizers for waiving their respective registrations fees.

References

Althaus, L. G., Serenelli, A. M., Panei, J. A., et al. 2005, A&A, 435, 631
Althaus, L. G., Miller Bertolami, M. M., Córsico, A. H., García-Berro, E., & Gil-Pons, P. 2005, A&A, 440, L1
Althaus, L. G., Córsico, A. H., Isern, J., & García-Berro, E. 2010, A&A Rev., 18, 471
Bergeron, P., Wesemael, F., Dufour, P., et al. 2011, ApJ, 737, 28
Bloecker, T. 1995, A&A, 299, 755
Blöcker, T. 2001, Ap&SS, 275, 1
Bradley, P. A. 2001, 12th European Workshop on White Dwarfs, 226, 307
Castanheira, B. G., & Kepler, S. O. 2009, MNRAS, 396, 1709
D'Antona, F., & Mazzitelli, I. 1990, ARA&A, 28, 139
Iben, I., Jr. 1984, ApJ, 277, 333
Iben, I., Jr., & MacDonald, J. 1986, ApJ, 301, 164
Iben, I., Jr., & MacDonald, J. 1995, White Dwarfs, 443, 48
Miller Bertolami, M. M., Alhaus, L. G., & Córsico, A. H. 2005, BAAA, 48, 185
Miller Bertolami, M. M., Althaus, L. G., Serenelli, A. M., & Panei, J. A. 2006, A&A, 449, 313
Miller Bertolami, M. M., & Althaus, L. G. 2007, MNRAS, 380, 763
Miller Bertolami, M. M., Althaus, L. G., Olano, C., & Jiménez, N. 2011, MNRAS, 415, 1396
Miller Bertolami, M. M. 2016, A&A, 588, A25
Paczyński, B. 1971, Acta Astron., 21, 417
Romero, A. D., Córsico, A. H., Althaus, L. G., et al. 2012, MNRAS, 420, 1462
Schoenberner, D. 1979, A&A, 79, 108
Schoenberner, D. 1987, Late Stages of Stellar Evolution, 132, 337
Tremblay, P.-E., & Bergeron, P. 2008, ApJ, 672, 1144-1152
Werner, K., & Herwig, F. 2006, PASP, 118, 183